arXiv:1511.06221v2 [math.NT] 21 Nov 2015

Preprint

PROOF OF SOME CONGRUENCES CONJECTURED BY GUO AND LIU

GUO-SHUAI MAO

Abstract. Let n and r be positive integers and p a prime. Define the numbers $S_n^{(r)}$ and $T_n^{(r)}$ by

$$S_n^{(r)}=\sum_{k=0}^{n}\binom{n}{k}^2\binom{2k}{k}(2k+1)^r\,,\ T_n^{(r)}=\sum_{k=0}^{n}\binom{n}{k}^2\binom{2k}{k}(2k+1)^r(-1)^k.$$

In this paper we first prove the following congruences:

$$\sum_{k=0}^{n-1}S_k^{(2r)}\equiv 0\ (\mathrm{mod}\ n^2),$$

$$\sum_{k=0}^{n-1}T_k^{(2r)}\equiv 0\ (\mathrm{mod}\ n^2)$$

and

$$\sum_{k=0}^{p-1}T_k^{(2)}\equiv\frac{p^2}{2}\left(5-3(\frac{p}{5})\right)\ (\mathrm{mod}\ p^3).$$

We also show that there exist integers a_{2r-1} and b_r, independent of n, such that

$$a_{2r-1}\sum_{k=0}^{n-1}S_k^{(2r-1)}\equiv 0\ (\mathrm{mod}\ n^2),$$

$$b_r\sum_{k=0}^{n-1}kS_k^{(r)}\equiv 0\ (\mathrm{mod}\ n^2).$$

This confirms several recent conjectures of Guo and Liu.

1. Introduction

In [Su1],Z.-W. Sun defined two new kinds of numbers,

$$S_n=\sum_{k=0}^{n}\binom{n}{k}^2\binom{2k}{k}(2k+1)\ (n=0,1,2,\ldots)$$

Key words and phrases. Central binomial coefficients, congruences, Bernoulli numbers, Zeilberger algorithm.

2010 *Mathematics Subject Classification.* 11B65, 11B68, 05A10, 11A07.

This research was supported by the Natural Science Foundation (Grant No. 11171140) of China.

and

$$R_n = \sum_{k=0}^{n} \binom{n}{k}\binom{n+k}{k}\frac{1}{2k-1}\ (n = 0, 1, 2, \ldots),$$

he showed a lot of congruences involving these two numbers, like

$$\frac{1}{n^2}\sum_{k=0}^{n-1} S_k = \sum_{k=0}^{n-1}\binom{n-1}{k}^2 C_k \in \mathbb{Z},$$

where $C_k = \frac{1}{k+1}\binom{2k}{k}$ is the k-th Catalan number, and

$$\frac{1}{n}\sum_{k=0}^{n-1} S_k(x) \in \mathbb{Z}[x].$$

And he gave lots of conjectures about these two numbers, such as

[Su1, Conjecture 5.4]. For every $n = 1, 2, 3\ldots$, we have

$$\frac{3}{n}\sum_{k=0}^{n-1} R_k^2 \in \mathbb{Z} \text{ and } \frac{1}{n}\sum_{k=0}^{n-1}(2k+1)R_k^2 \in \mathbb{Z}.$$

Conjecture 1.1. [Su1, Conjecture 5.5,5.6] *For any positive integer n, we have*

$$\frac{4}{n^2}\sum_{k=0}^{n-1} kS_k \in \mathbb{Z}, \frac{1}{n^2}\sum_{k=0}^{n-1} S_k^{(2)} \in \mathbb{Z}.$$

Guo and Liu proved some conjectures(contain Conjecture 1.1) of professor Z.W. Sun about S_n and other two numbers similar to S_n in [GL], they also defined two numbers

$$S_n^{(r)} = \sum_{k=0}^{n}\binom{n}{k}^2\binom{2k}{k}(2k+1)^r\ , T_n^{(r)} = \sum_{k=0}^{n}\binom{n}{k}^2\binom{2k}{k}(2k+1)^r(-1)^k,$$

which extend S_n and S_n^- in [Su1], note that $T_n^{(2)}$ is S_n^- introduced by Sun [Su1, Conjecture 5.6], Guo and Liu also gave few conjectures about $S_n^{(r)}$ and $T_n^{(r)}$ at the end of [GL], for example:

Conjecture 1.2. [GL, Conjecture 5.1] *Let n and r be positive integers and p a prime. Then*

$$\sum_{k=0}^{n-1} S_k^{(2r)} \equiv 0\ (\mathrm{mod}\ n^2),$$

$$\sum_{k=0}^{n-1} T_k^{(2r)} \equiv 0\ (\mathrm{mod}\ n^2),$$

$$\sum_{k=0}^{p-1} T_k^{(2)} \equiv \frac{p^2}{2}\left(5-3\left(\frac{p}{5}\right)\right) \pmod{p^3}.$$

Conjecture 1.3. [GL, Conjecture 5.2] *Let n and r be positive integers. Then there exist integers a_{2r-1} and b_r, independent of n, such that*

$$a_{2r-1}\sum_{k=0}^{n-1} S_k^{(2r-1)} \equiv 0 \pmod{n^2},$$

$$b_r\sum_{k=0}^{n-1} kS_k^{(r)} \equiv 0 \pmod{n^2}.$$

Conjecture 1.4. [GL, Conjecture 5.3] *Let*

$$a_3=3, a_5=15, a_7=21, a_9=15, a_{11}=33, a_{13}=1365, a_{15}=3.$$

$$b_2=12, b_3=4, b_4=60, b_5=20, b_6=84, b_7=28, b_8=60, b_9=20.$$

Then the conjecture 1.3 holds.

In this paper we mainly prove Conjecture 1.2 and Conjecture 1.3 , and partially answer the Conjecture 1.4.

We know that the Euler numbers $\{E_n\}$ and Euler polynomials $\{E_n(x)\}$ are defined by

$$\frac{2e^t}{e^{2t}+1}=\sum_{n=0}^{\infty} E_n\frac{t^n}{n!} \ (|t|<\frac{\pi}{2}) \text{ and } \frac{2e^{xt}}{e^t+1}=\sum_{n=0}^{\infty} E_n(x)\frac{t^n}{n!} \ (|t|<\pi).$$

And we also have the property of Euler polynomials

$$E_n(x+y)=\sum_{k=0}^{n}\binom{n}{k}E_k(x)y^{n-k}. \tag{1.1}$$

Recall that the Bernoulli numbers given by

$$\frac{x}{e^x-1}=\sum_{n=0}^{\infty} B_n\frac{x^n}{n!} \ \ (0<|x|<2\pi).$$

Motivated by the above work, we mainly obtain the following results in this paper.

Theorem 1.1. *For any positive integers n and r, we have*

$$\sum_{k=0}^{n-1} S_k^{(2r)} \equiv 0 \pmod{n^2}. \tag{1.2}$$

This result can detrude one conjecture in [Su1, Conjecture 5.6], that is

$$\frac{1}{n^2}\sum_{k=0}^{n-1} S_k^+ \in \mathbb{Z},$$

which has been solved by Guo and Liu in [GL].

Theorem 1.2. *Let n and r be positive integers, we can obtain*

$$\sum_{k=0}^{n-1} T_k^{(2r)} \equiv 0 \pmod{n^2}. \tag{1.3}$$

This result also can push out one conjecture in [Su1, Conjecture 5.6], that is

$$\frac{1}{n^2}\sum_{k=0}^{n-1} S_k^- \in \mathbb{Z},$$

where $S_k^- = T_k^{(2)}$.

The above two are divisibility results, following we give a congruence result:

Theorem 1.3. *Let p be a prime, then we have*

$$\sum_{k=0}^{n-1} T_k^{(2)} \equiv \frac{p^2}{2}\left(5 - 3\left(\frac{5}{p}\right)\right) \pmod{p^3}. \tag{1.4}$$

The following theorem is more challenging than the above three Theorems, because it is more difficult when the index number of S_k is odd.

Theorem 1.4. *Let n and r be positive integers, then there exist integers a_{2r-1} and b_r, independent of n, such that*

$$a_{2r-1}\sum_{k=0}^{n-1} S_k^{(2r-1)} \equiv 0 \pmod{n^2}, \tag{1.5}$$

$$b_r\sum_{k=0}^{n-1} kS_k^{(r)} \equiv 0 \pmod{n^2}. \tag{1.6}$$

We are going to prove Theorem 1.1 in section 2, then prove Theorem 1.2 in section 3, and at last we will prove Theorem 1.3 and Theorem 1.4 in section 4 and section 5 respectively, Our proofs make use of some sophisticated combinatorial identities and the Zeilberger algorithm.

We give two lemmas here, which will be used to prove the above four Theorems.

Lemma 1.1. *For any nonnegative integers m and n, we have*

$$\sum_{k=0}^{n}\binom{x+k}{m}=\binom{n+x+1}{m+1}-\binom{x}{m+1}, \tag{1.7}$$

and

$$\sum_{k=0}^{n}\binom{n}{k}^2\binom{x+k}{2n}=\binom{x}{n}^2. \tag{1.8}$$

Remark 1.1. Both (1.5) and (1.6) can be found in [G, (1.48) and (6.30)].

Lemma 1.2. [MS, (Lemma 3.2)] *For any nonnegative integer n, we have*

$$\sum_{k=0}^{n}\binom{n}{k}^2\binom{x+k}{2n+1}=\frac{1}{(4n+2)\binom{2n}{n}}\sum_{k=0}^{n}(2x-3k)\binom{x}{k}^2\binom{2k}{k}. \tag{1.9}$$

The identity (1.9) is crucial to our proofs of Theorems 1.1-1.4.

2. Proof of Theorem 1.1

Lemma 2.1. *For any positive integer n and r, we have*

$$\sum_{k=0}^{n-1}(2k+1)^{2r-1}\equiv 0\pmod{n}.$$

Proof: We know that

$$\sum_{k=0}^{n-1}(2k+1)^{2r-1}=\sum_{j=0}^{n-1}(2n-2j-1)^{2r-1}\equiv-\sum_{k=0}^{n-1}(2k+1)^{2r-1}\pmod{2n},$$

then

$$2\sum_{k=0}^{n-1}(2k+1)^{2r-1}\equiv 0\pmod{2n},$$

so we can deduce that

$$\sum_{k=0}^{n-1}(2k+1)^{2r-1}\equiv 0\pmod{n}.$$

Thus we have done the proof of Lemma 2.1.

Remark 2.1. We can see that in the proof of this lemma, we need the index number of S_k to be odd, if it is even, we can not get the result like this, so when the index number of S_k is even, the case is more difficult.

Proof of Theorem 1.1: By Lemma 1.1 we can deduce that

$$\begin{aligned}\sum_{k=0}^{n-1}S_k^{(2r)}&=\sum_{k=0}^{n-1}\sum_{j=0}^{k}\binom{k}{j}^2\binom{2j}{j}(2j+1)^{2r}\\&=\sum_{j=0}^{n-1}\binom{2j}{j}(2j+1)^{2r}\sum_{k=j}^{n-1}\sum_{l=0}^{j}\binom{j}{l}^2\binom{k+l}{2j}\\&=\sum_{j=0}^{n-1}\binom{2j}{j}(2j+1)^{2r}\sum_{l=0}^{j}\binom{j}{l}^2\sum_{k=j}^{n-1}\binom{k+l}{2j}\\&\equiv\sum_{j=0}^{n-1}\binom{2j}{j}(2j+1)^{2r}\sum_{l=0}^{j}\binom{j}{l}^2\binom{n+l}{2j+1}.\end{aligned}$$

Then via Lemma 1.2 and Lemma 2.1 we can obtain

$$\begin{aligned}&\sum_{k=0}^{n-1}S_k^{(2r)}=\frac{1}{2}\sum_{j=0}^{n-1}(2j+1)^{2r-1}\sum_{l=0}^{j}(2n-3l)\binom{n}{l}^2\binom{2l}{l}\\=&n\sum_{j=0}^{n-1}(2j+1)^{2r-1}+\frac{1}{2}\sum_{j=1}^{n-1}(2j+1)^{2r-1}\sum_{l=1}^{j}(2n-3l)\binom{n}{l}^2\binom{2l}{l}\\=&n\sum_{j=0}^{n-1}(2j+1)^{2r-1}+n^2\sum_{l=1}^{n-1}(2n-3l)\binom{n-1}{l-1}^2\frac{\binom{2l-1}{l-1}}{l^2}\sum_{j=l}^{n-1}(2j+1)^{2r-1}\\\equiv&n^2\sum_{l=1}^{n-1}(2n-3l)\binom{n-1}{l-1}^2\frac{\binom{2l-1}{l-1}}{l^2}\sum_{j=l}^{n-1}(2j+1)^{2r-1}\pmod{n^2}.\end{aligned}$$

We know that

$$\frac{\binom{n-1}{l-1}(2n-3l)}{l}=2\binom{n}{l}-3\binom{n-1}{l-1}\in\mathbb{Z}\text{ and }\frac{n}{l}\binom{n-1}{l-1}=\binom{n}{l}\in\mathbb{Z},$$

so with Lemma 2.1 we have $\sum_{j=0}^{n-1}(2j+1)^{2r-1}\equiv 0\pmod n$, then

$$\frac{n^2\binom{n-1}{l-1}^2(2n-3l)}{l^2}\sum_{j=0}^{n-1}(2j+1)^{2r-1}\equiv 0\pmod{n^2},$$

and $\sum_{j=0}^{l-1}(2j+1)^{2r-1}\equiv 0\pmod l$, then

$$\frac{n^2\binom{n-1}{l-1}^2(2n-3l)}{l^2}\sum_{j=0}^{l-1}(2j+1)^{2r-1}\equiv 0\pmod{n^2}.$$

Hence

$$n^2(2n-3l)\binom{n-1}{l-1}^2\frac{\binom{2l-1}{l-1}}{l^2}\sum_{j=l}^{n-1}(2j+1)^{2r-1}\equiv 0\ (\mathrm{mod}\ n^2).$$

Therefore

$$\sum_{k=0}^{n-1}S_k^{(2r)}\equiv 0\ (\mathrm{mod}\ n^2).$$

So we finish the proof of Theorem 1.1. □

3. Proof of Theorem 1.2

Lemma 3.1. *For any positive integer n and r ,we have*

$$\sum_{k=0}^{n-1}(2k+1)^{2r-1}(-1)^k\equiv 0\ (\mathrm{mod}\ n).$$

Proof: By $E_n(x)+E_n(x+1)=2x^n$, $E_n(\frac{1}{2})=\frac{E_n}{2^n}$ and (1.1), we have

$$\begin{aligned}
&\sum_{k=0}^{n-1}(-1)^k(2k+1)^{2r-1}\\
=&2^{2r-2}\sum_{k=0}^{n-1}((-1)^kE_{2r-1}(k+\frac{1}{2})-(-1)^{k+1}E_{2r-1}(k+\frac{3}{2}))\\
=&2^{2r-2}(E_{2r-1}(\frac{1}{2})-(-1)^nE_{2r-1}(n+\frac{1}{2}))\\
=&\frac{E_{2r-1}}{2}-(-1)^n\frac{E_{2r-1}}{2}-(-1)^n2^{2r-2}\sum_{k=0}^{2r-2}\binom{2r-1}{k}\frac{E_k}{2^k}n^{2r-1-k}\\
\equiv&0\ (\mathrm{mod}\ n).
\end{aligned}$$

(Note that $E_{2n-1}=0$ and $E_n\in\mathbb{Z}$ for all $n\geq 1$, $\frac{2^{2r-2}}{2^k}\in\mathbb{Z}$ for all $k=0,1,\cdots,2r-2$.)

So we complete the proof of Lemma 3.1.

Remark 3.1. We know that $E_k=0$ when k is odd and $E_n\in\mathbb{Z}$ for all $n\geq 1$, so we can prove this lemma by the above method. If the index number of $2k+1$ is not $2r-1$, it is an even number $2r$, then we can not prove it like this.

Proof of Theorem 1.2: By Lemma 1.1 we can deduce that

$$\begin{aligned}\sum_{k=0}^{n-1}T_k^{(2r)}&=\sum_{k=0}^{n-1}\sum_{j=0}^{k}\binom{k}{j}^2\binom{2j}{j}(2j+1)^{2r}(-1)^j\\&=\sum_{j=0}^{n-1}\binom{2j}{j}(2j+1)^{2r}(-1)^j\sum_{k=j}^{n-1}\sum_{l=0}^{j}\binom{j}{l}^2\binom{k+l}{2j}\\&=\sum_{j=0}^{n-1}\binom{2j}{j}(2j+1)^{2r}(-1)^j\sum_{l=0}^{j}\binom{j}{l}^2\sum_{k=j}^{n-1}\binom{k+l}{2j}\\&\equiv\sum_{j=0}^{n-1}\binom{2j}{j}(2j+1)^{2r}(-1)^j\sum_{l=0}^{j}\binom{j}{l}^2\binom{n+l}{2j+1}.\end{aligned}$$

Then via Lemma 1.2 and Lemma 3.1 we can obtain

$$\begin{aligned}&\sum_{k=0}^{n-1}T_k^{(2r)}=\frac{1}{2}\sum_{j=0}^{n-1}(2j+1)^{2r-1}(-1)^j\sum_{l=0}^{j}(2n-3l)\binom{n}{l}^2\binom{2l}{l}\\=&n\sum_{j=0}^{n-1}(2j+1)^{2r-1}(-1)^j+\frac{1}{2}\sum_{j=1}^{n-1}(2j+1)^{2r-1}(-1)^j\sum_{l=1}^{j}(2n-3l)\binom{n}{l}^2\binom{2l}{l}\\=&n\sum_{j=0}^{n-1}(2j+1)^{2r-1}(-1)^j+n^2\sum_{l=1}^{n-1}(2n-3l)\frac{\binom{n-1}{l-1}^2\binom{2l-1}{l-1}}{l^2}\sum_{j=l}^{n-1}(2j+1)^{2r-1}(-1)^j\\\equiv&n^2\sum_{l=1}^{n-1}(2n-3l)\binom{n-1}{l-1}^2\frac{\binom{2l-1}{l-1}}{l^2}\sum_{j=l}^{n-1}(2j+1)^{2r-1}(-1)^j\pmod{n^2}.\end{aligned}$$

We know that

$$\frac{\binom{n-1}{l-1}(2n-3l)}{l}=2\binom{n}{l}-3\binom{n-1}{l-1}\in\mathbb{Z}\text{ and }\frac{n}{l}\binom{n-1}{l-1}=\binom{n}{l}\in\mathbb{Z},$$

so with Lemma 3.1 we have $\sum_{j=0}^{n-1}(2j+1)^{2r-1}(-1)^j\equiv 0\pmod n$, then

$$\frac{n^2\binom{n-1}{l-1}^2(2n-3l)}{l^2}\sum_{j=0}^{n-1}(2j+1)^{2r-1}(-1)^j\equiv 0\pmod{n^2},$$

and $\sum_{j=0}^{l-1}(2j+1)^{2r-1}(-1)^j\equiv 0\pmod l$, then

$$\frac{n^2\binom{n-1}{l-1}^2(2n-3l)}{l^2}\sum_{j=0}^{l-1}(2j+1)^{2r-1}(-1)^j\equiv 0\pmod{n^2}.$$

So we can get that

$$n^2(2n-3l)\binom{n-1}{l-1}^2\frac{\binom{2l-1}{l-1}}{l^2}\sum_{j=l}^{n-1}(2j+1)^{2r-1}(-1)^j\equiv 0\ (\mathrm{mod}\ n^2).$$

Then we have completed the proof of Theorem 1.2. □

4. Proof of Theorem 1.3

Proof of Theorem 1.3: By Lemma 1.1 we can deduce that

$$\begin{aligned}\sum_{k=0}^{p-1}T_k^{(2)}&=\sum_{k=0}^{p-1}\sum_{j=0}^{k}\binom{k}{j}^2\binom{2j}{j}(2j+1)^2(-1)^j\\&=\sum_{j=0}^{p-1}\binom{2j}{j}(2j+1)^2(-1)^j\sum_{k=j}^{p-1}\sum_{l=0}^{j}\binom{j}{l}^2\binom{k+l}{2j}\\&=\sum_{j=0}^{p-1}\binom{2j}{j}(2j+1)^2(-1)^j\sum_{l=0}^{j}\binom{j}{l}^2\sum_{k=j}^{p-1}\binom{k+l}{2j}\\&=\sum_{j=0}^{p-1}\binom{2j}{j}(2j+1)^2(-1)^j\sum_{l=0}^{j}\binom{j}{l}^2\binom{p+l}{2j+1}.\end{aligned}$$

Then via Lemma 1.2 we can obtain

$$\begin{aligned}&\sum_{k=0}^{p-1}T_k^{(2)}=\frac{1}{2}\sum_{j=0}^{p-1}(2j+1)(-1)^j\sum_{l=0}^{j}(2p-3l)\binom{p}{l}^2\binom{2l}{l}\\=&p\sum_{j=0}^{p-1}(2j+1)(-1)^j+\frac{1}{2}\sum_{j=1}^{p-1}(2j+1)(-1)^j\sum_{l=1}^{j}(2p-3l)\binom{p}{l}^2\binom{2l}{l}\\=&p\sum_{j=0}^{p-1}(2j+1)(-1)^j+\frac{p^2}{2}\sum_{l=1}^{p-1}(2p-3l)\binom{p-1}{l-1}^2\frac{\binom{2l}{l}}{l^2}\sum_{j=l}^{p-1}(2j+1)(-1)^j\\\equiv&p^2-\frac{3p^2}{2}\sum_{l=1}^{p-1}\frac{\binom{2l}{l}}{l}\binom{p-1}{l-1}^2\sum_{j=l}^{p-1}(2j+1)(-1)^j\\\equiv&p^2-\frac{3}{2}p^2\sum_{l=1}^{p-1}\binom{2l}{l}(-1)^l\ (\mathrm{mod}\ p^3),\end{aligned}$$

(Note that $\sum_{j=0}^{p-1}(2j+1)(-1)^j=p$, $\sum_{j=l}^{p-1}(2j+1)(-1)^j=p+(-1)^l l$ and $\binom{p-1}{l-1}^2=\Pi_{j=1}^{l-1}(\frac{p}{j}-1)^2\equiv 1\ (\mathrm{mod}\ p)$).

We know that $\binom{2l}{l} = \binom{-\frac{1}{2}}{l}(-4)^l \equiv \binom{\frac{p-1}{2}}{l}(-4)^l \pmod p$, so we have

$$\begin{aligned}\sum_{k=0}^{p-1} T_k^{(2)} \equiv & p^2 - \frac{3}{2}p^2 \sum_{l=1}^{(p-1)/2} \binom{\frac{p-1}{2}}{l} 4^l \\ \equiv & p^2 - \frac{3}{2}p^2(5^{(p-1)/2} - 1) \equiv \frac{p^2}{2}\left(5 - 3\left(\frac{5}{p}\right)\right) \pmod{p^3}.\end{aligned}$$

Note that $\left(\frac{5}{p}\right) = \left(\frac{p}{5}\right)$, so we finish the proof of Theorem 1.3. □

5. Proof of Theorem 1.4

In [KM] we know $S_m(n) = 1^m + 2^m + \cdots + (n-1)^m$ and the m-th Bernoulli number can be written as $B_m = U_m/V_m$ where $(U_m, V_m) = 1$ and $V_m > 0$.

Lemma 5.1. [KM, Proposition 15.2.2] *If integer $m \geq 2$ is even, then for all $n \geq 1$ we have*

$$V_m S_m(n) \equiv nU_m \pmod{n^2}.$$

Lemma 5.2. *Let n and r be positive integers and $r > 1$, then we have*

$$V_{2r-2} \sum_{j=0}^{n-1} (2j+1)^{2r-2} \equiv 0 \pmod{2n}.$$

Proof: We know $\sum_{j=0}^{n-1}(2j+1)^{2r-2} = S_{2r-2}(2n) - 2^{2r-2}S_{2r-2}(n)$, and by Lemma 5.1 we have

$$V_{2r-2}S_{2r-2}(2n) \equiv 2nU_{2r-2} \equiv 0 \pmod{2n}$$

and

$$V_{2r-2}S_{2r-2}(n) \equiv nU_{2r-2} \equiv 0 \pmod{n}$$

for all $r > 1$, so for all positive integer $r > 1$ we have

$$V_{2r-2} \sum_{j=0}^{n-1} (2j+1)^{2r-2} \equiv 0 \pmod{2n}.$$

Therefore we have done the proof of Lemma 5.2.
Proof of (1.5): For $r = 1$ we know $\sum_{k=0}^{n-1} S_k \equiv 0 \pmod{n^2}$, which is given by Z.W.Sun, so we just need to show it for all $r > 1$. By Lemma

1.1 we can deduce that

$$\begin{aligned}\sum_{k=0}^{n-1}S_k^{(2r-1)}&=\sum_{k=0}^{n-1}\sum_{j=0}^{k}\binom{k}{j}^2\binom{2j}{j}(2j+1)^{2r-1}\\&=\sum_{j=0}^{n-1}\binom{2j}{j}(2j+1)^{2r-1}\sum_{k=j}^{n-1}\sum_{l=0}^{j}\binom{j}{l}^2\binom{k+l}{2j}\\&=\sum_{j=0}^{n-1}\binom{2j}{j}(2j+1)^{2r-1}\sum_{l=0}^{j}\binom{j}{l}^2\sum_{k=j}^{n-1}\binom{k+l}{2j}\\&=\sum_{j=0}^{n-1}\binom{2j}{j}(2j+1)^{2r-1}\sum_{l=0}^{j}\binom{j}{l}^2\binom{n+l}{2j+1}.\end{aligned}$$

Then via Lemma 1.2 and Lemma 5.2 we can obtain

$$\begin{aligned}V_{2r-2}\sum_{k=0}^{n-1}S_k^{(2r-1)}&=\frac{V_{2r-2}}{2}\sum_{j=0}^{n-1}(2j+1)^{2r-2}\sum_{l=0}^{j}(2n-3l)\binom{n}{l}^2\binom{2l}{l}\\&=V_{2r-2}n\sum_{j=0}^{n-1}(2j+1)^{2r-2}+\frac{V_{2r-2}}{2}\sum_{j=1}^{n-1}(2j+1)^{2r-2}\sum_{l=1}^{j}(2n-3l)\binom{n}{l}^2\binom{2l}{l}\\&=V_{2r-2}n\sum_{j=0}^{n-1}(2j+1)^{2r-2}+n^2\sum_{l=1}^{n-1}\frac{V_{2r-2}(2n-3l)\binom{n-1}{l-1}^2\binom{2l-1}{l-1}}{l^2}\sum_{j=l}^{n-1}(2j+1)^{2r-2}\\&\equiv V_{2r-2}n^2\sum_{l=1}^{n-1}(2n-3l)\binom{n-1}{l-1}^2\frac{\binom{2l-1}{l-1}}{l^2}\sum_{j=l}^{n-1}(2j+1)^{2r-2}\pmod{2n^2}.\end{aligned}$$

We know $\frac{\binom{n-1}{l-1}(2n-3l)}{l}=2\binom{n}{l}-3\binom{n-1}{l-1}\in\mathbb{Z}$ and $\frac{n\binom{n-1}{l-1}}{l}=\binom{n}{l}\in\mathbb{Z}$, so with Lemma 5.2 we have

$$\frac{V_{2r-2}n^2\binom{n-1}{l-1}^2(2n-3l)}{l^2}\sum_{j=0}^{n-1}(2j+1)^{2r-2}\equiv 0\pmod{2n^2},$$

and

$$\frac{V_{2r-2}n^2\binom{n-1}{l-1}^2(2n-3l)}{l^2}\sum_{j=0}^{l-1}(2j+1)^{2r-2}\equiv 0\pmod{2n^2}.$$

So we can get that

$$V_{2r-2}n^2(2n-3l)\binom{n-1}{l-1}^2\frac{\binom{2l-1}{l-1}}{l^2}\sum_{j=l}^{n-1}(2j+1)^{2r-2}\equiv 0\pmod{2n^2}.$$

Therefore

$$V_{2r-2}\sum_{k=0}^{n-1}S_k^{(2r-1)}\equiv 0\ (\mathrm{mod}\ 2n^2),$$

so we just need to take

$$\begin{cases} a_{2r-1}=1 & \text{if r=1,}\\ a_{2r-1}=\dfrac{1}{2}V_{2r-2} & \text{if r>1.}\end{cases}$$

Then we have finished the proof of (1.5).

Lemma 5.3. *For any positive integer n and $0\le j<n$ we have*

$$(2j+1)\binom{2j}{j}\sum_{k=j}^{n-1}(2k-j+1)\binom{k}{j}^2\equiv 0\ (\mathrm{mod}\ n^2).$$

Proof: Set $u_j=(2j+1)\binom{2j}{j}\sum_{k=j}^{n-1}(2k-j+1)\binom{k}{j}^2$, then by Zeilberger we can get the recurrence

$$u_j=\frac{(1+2j)n^2\binom{2j}{j}\binom{n-1}{j}^2}{j+1}$$

for all $0\le j<n-1$, while $\frac{\binom{2j}{j}}{j+1}\in\mathbb{Z}$, so we have $u_j\equiv 0\ (\mathrm{mod}\ n^2)$ for all $0\le j<n-1$, and we know

$$u_{n-1}=(2n-1)\binom{2n-2}{n-1}n=n^2\binom{2n-1}{n-1}\equiv 0\ (\mathrm{mod}\ n^2),$$

so $u_j\equiv 0\ (\mathrm{mod}\ n^2)$ for all $0\le j<n$, therefore we finish the proof of Lemma 5.3.

Proof of (1.6): First we know that

$$\begin{aligned}&\sum_{k=0}^{n-1}(4kS_k^{(r)}-S_k^{(r+1)}+3S_k^{(r)})\\ =&\sum_{k=0}^{n-1}\sum_{j=0}^{k}(4k-2j+2)\binom{k}{j}^2\binom{2j}{j}(2j+1)^r\\ =&\sum_{j=0}^{n-1}\binom{2j}{j}(2j+1)^r\sum_{k=j}^{n-1}(4k-2j+2)\binom{k}{j}^2=2\sum_{j=0}^{n-1}(2j+1)^{r-1}u_j\end{aligned}$$

then by Lemma 5.3 we can deduce that

$$\sum_{k=0}^{n-1}(4kS_k^{(r)}-S_k^{(r+1)}+3S_k^{(r)})\equiv 0\ (\mathrm{mod}\ 2n^2),$$

we know when $r = 1$, $\sum_{k=0}^{n-1} 4kS_k^{(r)} \equiv 0 \pmod{n^2}$, which is given by Guo and Liu [GL]. so we just need to show it for all $r > 1$.

Case 1. If $2|r$, then by (1.2) we have $V_r \sum_{k=0}^{n-1} S_k^{(r)} \equiv 0 \pmod{2n^2}$ since 6 always divides the denominator of B_r [KM, Page 233,Corollary], and with (1.5) we have $V_r \sum_{k=0}^{n-1} S_k^{(r+1)} \equiv 0 \pmod{2n^2}$, so we can get

$$V_r \sum_{k=0}^{n-1} 4kS_k^{(r)} \equiv 0 \pmod{2n^2}.$$

Case 2. If $2 \nmid r$ and $r > 1$, then by (1.2) we have $V_{r-1} \sum_{k=0}^{n-1} S_k^{(r+1)} \equiv 0 \pmod{2n^2}$ since 6 always divides the denominator of B_{r-1} [KM, Page 233,Corollary], and with (1.5) we know $V_{r-1} \sum_{k=0}^{n-1} S_k^{(r)} \equiv 0 \pmod{2n^2}$, thus we have

$$V_{r-1} \sum_{k=0}^{n-1} 4kS_k^{(r)} \equiv 0 \pmod{2n^2}.$$

So combining Case 1 and Case 2 we finally get that there exist integers b_r such that

$$b_r \sum_{k=0}^{n-1} kS_k^{(r)} \equiv 0 \pmod{n^2},$$

we just need to take

$$\begin{cases} b_r = 2V_{r-1} & \text{if r} \equiv 1 \pmod 2, \\ b_r = 2V_r & \text{if r} \equiv 0 \pmod 2. \end{cases}$$

Therefore we complete the proof of (1.6). □

Corollary 5.1. *We can deduce some special value of a_{2r-1}, like*

$$a_3 = 3, a_5 = 15, a_7 = 21, a_9 = 15, a_{11} = 33, a_{13} = 1365, a_{15} = 3.$$

Proof: We know

$$B_2 = \frac{1}{6}, B_4 = -\frac{1}{30}, B_6 = \frac{1}{42}, B_8 = -\frac{1}{30},$$

$$B_{10} = \frac{5}{66}, B_{12} = -\frac{691}{2730}, B_{14} = \frac{7}{6},$$

so we have

$$V_2 = 6, V_4 = 30, V_6 = 42, V_8 = 30, V_{10} = 66, V_{12} = 2730, V_{14} = 6,$$

then by (1.5) we can deduce that

$$a_3 = 3, a_5 = 15, a_7 = 21, a_9 = 15, a_{11} = 33, a_{13} = 1365, a_{15} = 3.$$

Corollary 5.2. *We can deduce some special value of b_r, such as*

$$b_2 = b_3 = 12, b_4 = b_5 = 60, b_6 = b_7 = 84, b_8 = b_9 = 60,$$

and

$$b_{10} = b_{11} = 132, b_{12} = b_{13} = 5460, b_{14} = b_{15} = 12.$$

Proof: We know $b_{2r+1} = b_{2r}$ in the proof of (1.6), and by (1.6) and

$$V_2 = 6, V_4 = 30, V_6 = 42, V_8 = 30, V_{10} = 66, V_{12} = 2730, V_{14} = 6.$$

we have

$$b_2 = 12, b_4 = 60, b_6 = 84, b_8 = 60, b_{10} = 132, b_{12} = 5460, b_{14} = 12$$

Thus

$$b_2 = b_3 = 12, b_4 = b_5 = 60, b_6 = b_7 = 84, b_8 = b_9 = 60,$$

and

$$b_{10} = b_{11} = 132, b_{12} = b_{13} = 5460, b_{14} = b_{15} = 12.$$

Remark 5.1. The corollary 5.1 and 5.2 partially answer the conjecture 5.3 of Guo and Liu in [GL]. □

Acknowledgment. The author would like to thank professors Z.-W. Sun and Hao Pan for helpful comments.

References

[G] H.W. Gould, *Combinatorial Identities*, Morgantown Printing and Binding Co., 1972.

[GL] V.J.W. Guo and J.-C. Liu, *Proof of some conjectures of Z.-W.Sun on the divisibility of certain double-sums*, to appear.

[KM] K. Ireland and M. Rosen, *A Classical Introduction to Modern Number Theory*, 2nd Editor, Grad. Texts in Math., Vol.84, Springer, New York, 1990.

[MS] G.-S. Mao and Z.W. Sun, *Two congruences involving haemonic numbers with applications*, to appear.

[Su1] Z.W. Sun, *Two new kinds of numbers and related divisibility results*, Preprint, `arXiv:1408.5381`, 2014.

(Guo-Shuai Mao) Department of Mathematics, Nanjing University, Nanjing 210093, People's Republic of China

E-mail address: `mg1421007@smail.nju.edu.cn`